

Question 2.

We will denote by H ($H = \text{const}$) the height of the tube above the mercury level in the pan, and the height of the mercury column in the tube by h_i . Under conditions of mechanical equilibrium the hydrogen pressure in the tube is:

$$(2.1) \quad P_{H_2} = P_{air} - \rho g h_i,$$

where ρ is the density of mercury at temperature t_i :

$$(2.2) \quad \rho = \rho_0(1 - \beta t)$$

The index i enumerates different stages undergone by the system, ρ_0 is the density of mercury at $t_0 = 0^\circ\text{C}$, or $T_0 = 273\text{ K}$, and β its coefficient of expansion. The volume of the hydrogen is given by:

$$(2.3) \quad V_i = S(H - h_i).$$

Now we can write down the equations of state for hydrogen at points 0, 1, 2, and 3 of the PV diagram (see Fig. 2):

$$(2.4) \quad (P_0 - \rho_0 g h_0)S(H - h_0) = \frac{m}{M} RT_0;$$

$$(2.5) \quad (P_1 - \rho_0 g h_1)S(H - h_1) = \frac{m}{M} RT_0;$$

$$(2.6) \quad (P_2 - \rho_1 g h_2)S(H - h_2) = \frac{m}{M} RT_2,$$

where $P_2 = \frac{P_1 T_2}{T_0}$, $\rho_1 = \frac{\rho_0}{1 + \beta(T_2 - T_0)} \approx \rho_0[1 - \beta(T_2 - T_0)]$ since the process 1–3 is isochoric, and:

$$(2.7) \quad (P_2 - \rho_2 g h_3)S(H - h_3) = \frac{m}{M} RT_3$$

where $\rho_2 \approx \rho_0[1 - \beta(T_3 - T_0)]$, $T_3 = T_2 \frac{V_3}{V_2} = T_2 \frac{H - h_3}{H - h_2}$ for the isobaric process 2–3.




Fig. 2

After a good deal of algebra the above system of equations can be solved for the unknown quantities, an exercise, which is left to the reader. The numerical answers, however, will be given for reference:

$H \approx 1.3 \text{ m};$
 $m \approx 2.11 \times 10^{-6} \text{ kg};$
 $T_2 \approx 364 \text{ K};$
 $P_2 \approx 1.067 \times 10^5 \text{ Pa};$
 $T_3 \approx 546 \text{ K};$
 $P_2 \approx 4.8 \times 10^4 \text{ Pa}.$