

Solution of problem 3:

a) Floating condition:

The total mass of the balloon, consisting of the mass of the envelope m_H and the mass of the air quantity of temperature ϑ_2 must equal the mass of the displaced air quantity with temperature $\vartheta_1 = 20^\circ\text{C}$.

$$V_B \cdot \rho_2 + m_H = V_B \cdot \rho_1$$

$$\rho_2 = \rho_1 - \frac{m_H}{V_B} \tag{1}$$

Then the temperature may be obtained from

$$\frac{\rho_1}{\rho_2} = \frac{T_2}{T_1},$$

$$T_2 = \frac{\rho_1}{\rho_2} \cdot T_1 = 341.53 \text{ K} = 68.38 \text{ }^\circ\text{C} \quad (2)$$

- b) The force F_B acting on the rope is the difference between the buoyant force F_A and the weight force F_G :

$$F_B = V_B \cdot \rho_1 \cdot g - (V_B \cdot \rho_3 + m_H) \cdot g \quad (3)$$

It follows with $\rho_3 \cdot T_3 = \rho_1 \cdot T_1$

$$F_B = V_B \cdot \rho_1 \cdot g \cdot \left(1 - \frac{T_1}{T_3}\right) - m_H \cdot g = 1,21 \text{ N} \quad (4)$$

- c) The balloon rises to the height h , where the density of the external air ρ_h has the same value as the effective density ρ_{eff} , which is evaluated from the mass of the air of temperature $\vartheta_3 = 110 \text{ }^\circ\text{C}$ (inside the balloon) and the mass of the envelope m_H :

$$\rho_{\text{eff}} = \frac{m_2}{V_B} = \frac{\rho_3 \cdot V_B + m_H}{V_B} = \rho_h = \rho_1 \cdot e^{\frac{\rho_1 \cdot g \cdot h}{\rho_0}} \quad (5)$$

Resolving eq. (5) for h gives: $h = \frac{p_0}{\rho_1 \cdot g} \cdot \ln \frac{\rho_1}{\rho_{\text{eff}}} = 843 \text{ m} \quad (6).$

- d) For *small* height differences (10 m in comparison to 843 m) the exponential pressure drop (or density drop respectively) with height can be approximated by a linear function of height. Therefore the driving force is proportional to the elongation out of the equilibrium position.

This is the condition in which harmonic oscillations result, which of course are damped by the air resistance.